Physical & Biological Sciences #672

Beyond Neural Activity: The Emergence of Consciousness through Light Refraction, Pressure Gradients, and Modulated Entropy

A Framework for Conserved Entropic-Photonic Symmetry

Angel Motto | Independent Researcher | Brandeis University

We propose that consciousness emerges when **photonic coherence**, **pressure gradients**, and **entropy regulation** interact in **cascades** that scale coherent patterns across the nervous system.

PROBLEM & EVIDENCE

The Explanatory Gap


Despite substantial progress in identifying neural correlates and large-scale information-processing mechanisms associated with consciousness, no single framework currently explains how conscious states emerge and reorganize across multiple biological scales.

Existing theories emphasize different levels of neural organization, while the possible contributions of intracellular physical processes, entropy regulation, and cross-scale dynamics remain comparatively unresolved.

This leaves an open mechanistic question: how does the nervous system maintain both stability and the capacity for rapid reorganization as conscious states evolve?

Empirical Foundations


Independent Empirical Streams



Independent findings support constituent phenomena and dynamical principles relevant to the framework; their proposed cross-scale coupling remains unresolved.

PROPOSED MECHANISM

Entropy-Window State Dynamics: Temporal Structure of State-Space Evolution



- **Accessible state space:** Neural dynamics occupy a bounded space of possible states shaped by ongoing physical, contextual, and biological constraints.
- **Constraint → proposed reduction:** Progressive constraint narrows accessible trajectories toward transient convergence. The framework proposes that this transition may culminate in a discrete quantum reduction event, potentially linking the model to Orch OR.
- **Probabilistic re-emergence:** Reduction is not treated as an endpoint. The realized state establishes new conditions from which a transformed distribution of possible states emerges.
- **Recursive evolution:** These translated constraints shape the next entropy window, producing a recurrent sequence of possibility → constraint → reduction → re-emergence → reconfiguration across time.


Quantum reduction is proposed as a candidate mechanism for transient selection; its role in neuronal dynamics remains theoretical.

- **Relation to Orch OR:** Orch OR proposes coherent and entangled quantum processes within neuronal microtubules culminating in objective reduction. The present framework extends this possibility by proposing that nested quantum correlations may persist across recursive state translations, preserving relational organization while constraining subsequent entropy-window accessibility.

Persistent microtubular entanglement, cross-scale propagation of quantum correlations, and symmetry preservation across neural state transitions remain theoretical.

- **Optical conditions:** Refractive boundaries may permit localized light confinement within or around microtubules under suitable conditions.
- **Proposed physical coupling:** Photonic activity may interact with the tubulin lattice and surrounding fluid, potentially influencing structural and pressure dynamics.
- **Neuronal translation:** These proposed interactions could alter ionic conditions, membrane excitability, and neuronal response probabilities.
- **Entropy-window modulation:** Changes in neuronal and network dynamics may reshape which neural states remain dynamically accessible.
- **Recursive reconfiguration:** Each realized state changes the conditions governing what can emerge next.

Microtubular optical confinement, pressure-mediated coupling, and their proposed influence on neural-state dynamics remain to be experimentally established.




THEORETICAL IMPLICATIONS

Predictions of the Model

- Coupled physical dynamics: If the proposed optical–mechanical mechanism operates, changes in microtubule-associated optical signals should covary with measurable lattice or local fluid dynamics under controlled conditions.
- Sensitivity to boundary conditions: Manipulating the local optical or physicochemical environment should alter any proposed confinement or coupling effects in a reproducible, condition-dependent manner.
- Neuronal consequences: If the coupling is functionally relevant, perturbing it should produce measurable changes in neuronal excitability or response probabilities beyond effects attributable to established cellular pathways.
- State-space reconfiguration: If these cellular changes influence network dynamics, targeted perturbations should alter the accessibility or transition probabilities of neural states, rather than merely changing overall activity.

Falsification Criteria & Experimental Tests



Unified Symmetry of Awareness

Consciousness as the dynamic conservation of relational form